

On complex surfaces with 5 or 6 semistable singular fibers over \mathbb{P}^1 *

Sheng-Li Tan

Department of Mathematics, East China Normal University,
Shanghai 200062, P. R. of China
sltan@math.ecnu.edu.cn

Yuping Tu

Department of Mathematics, China Three Gorges University,
Yichang 443002, Hubei Province, P. R. of China
tuyu02@sina.com

Alexis G. Zamora

CIMAT, AP 402, C.P. 36000, Guanajuato, Gto., México
alexis@cimat.mx

Introduction

We denote by X a complex smooth projective surface, and by $f : X \rightarrow C$ a fibration over a curve C whose generic fiber is a curve of genus g . f is called *isotrivial* if all smooth fibers are isomorphic to a fixed curve. f is called *semistable* if all of the singular fibers are reduced nodal curves. If there is no (-1) -curve contained in the fibers, then we call f *relatively minimal*. The projection $X = F \times C \rightarrow C$ is called a trivial fibration. s is always the number of singular fibers of f .

In a well-known paper [3], Beauville proved that a non-isotrivial fibration $f : X \rightarrow \mathbb{P}^1$ admits at least 3 singular fibers, and if f is semistable, then $s \geq 4$. The first author proved in [11] that if f is a semistable fibration of genus $g \geq 2$, then it admits at least 5 singular fibers.

Beauville [4] gave a beautiful explicit classification of all semistable elliptic fibrations ($g = 1$) over \mathbb{P}^1 with 4 singular fibers. More precisely, there are exactly 6 non-isotrivial such families, and all of them are modular families of elliptic curves. A very interesting natural problem is to classify semistable fibrations over \mathbb{P}^1 of genus $g \geq 2$ with 5 singular fibers. In the survey ([13], §1 and Ex. 5.9), Viehweg considered this problem. The purpose of this note is to try to get some information on the structure of the surface X when f has 5 or 6 singular fibers. Our main results are the following two theorems.

Theorem 0.1. *Assume that $f : X \rightarrow \mathbb{P}^1$ is a non trivial semistable fibration of genus $g \geq 2$ with s singular fibers. Assume also that f is relatively minimal. Then we have:*

*Keywords: singular fiber, fibration, surface, canonical class inequality. The first author would like to thank the support of “DFG-Schwerpunktprogramm Globale Methoden in der Komplexen Geometrie” and the DFG-NSFC Chinese-German project “Komplexe Geometrie”. He is also supported by the 973 Project Foundation and the Foundation of EMC for Doctoral Program. The third author is partially supported by Conacyt Grant 41459-F.

- (1) If $s = 5$, then X is birationally rational or ruled.
- (2) If $s = 6$ and $g = 2, 3$ or 4 , then X is not of general type.
- (3) If $s = 6$, $g = 5$ and X is of general type, then the minimal model S of X satisfies

$$K_S^2 = 1, \quad p_g(S) = 2, \quad q(S) = 0.$$

The fibration f comes from a pencil on S with 5 simple base points (including infinitely near base points).

Thus if the Kodaira dimension of X is non negative, then $s \geq 6$. We have no example of case (3), and we conjecture that s is at least 7 when X is of general type. We shall give two examples to show that the two bounds are sharp.

- A) $s = 6$, $g = 3$, $K_X^2 = -4$, and X is birationally a K3 surface.
- B) $s = 7$, $g = 4$, $K_X^2 = -3$, $p_g(X) = 2$, $q(X) = 0$, and X is of general type containing 4 (-1) -curves.

Our next purpose is to give some inequalities for genus $g \geq 2$ fibrations over \mathbb{P}^1 (not necessarily semistable), which imply the lower bounds on s .

Theorem 0.2. *Let $f : X \rightarrow \mathbb{P}^1$ be a relatively minimal fibration of genus $g \geq 2$, and let $K_f = K_{X/\mathbb{P}^1} = K_X + 2F$ be its relative canonical divisor. Assume that f is not locally trivial. Then*

$$K_f^2 \geq \begin{cases} 4g - 4, & \text{if } \kappa(X) = -\infty, \\ 6g - 6, & \text{if } \kappa(X) = 0, \\ 6g - 5, & \text{if } \kappa(X) = 1, \\ 6g - 6 + \frac{1}{2} \left(K_S^2 + \sqrt{K_S^2} \sqrt{K_S^2 + 8g - 8} \right), & \text{if } \kappa(X) = 2, \end{cases}$$

where S is the unique minimal model of X when $\kappa(X) \geq 0$.

The bounds are optimal for infinitely many g . The example given in [11] satisfies $K_f^2 = 4(g - 1)$, $g = 2$ and $s = 5$. The example A) above satisfies $K_f^2 = 6(g - 1)$, $g = 3$ and $s = 6$, where X is a K3 surface. In Theorem 2.1, we will classify the fibrations $f : X \rightarrow \mathbb{P}^1$ with minimal K_f^2 according to its Kodaira dimension.

The proof of Theorem 0.1 is to use some inequalities, particularly the following strict canonical class inequality and its refinement for a non trivial semistable fibration $f : X \rightarrow C$ of genus $g \geq 2$ with $s \neq 0$ [11] (see also [9] for a differential geometric proof):

$$K_{X/C}^2 < (2g - 2)(2g(C) - 2 + s). \quad (0.1)$$

We use Reider's method to prove Theorem 0.2. Note that the strict canonical class inequality and the inequality $K_f^2 \geq 6(g - 1)$ (resp. $K_f^2 \geq 4(g - 1)$) in Theorem 0.2 imply that $s \geq 6$ (resp. 5).

1 Preliminaries

Let $f : X \rightarrow C$ be a fibration of genus $g \geq 2$, namely X (resp. C) is a nonsingular complex surface (resp. curve) and the generic fiber F of f is a nonsingular curve of genus g . We always assume that f is relatively minimal, i.e., there is no (-1) -curve contained in the fibers. f is called *semistable* if all of the singular fibers are reduced nodal curves.

Denote by $K_f = K_{X/C} = K_X - f^*K_C$ the relative canonical divisor of f , and by $\omega_{X/C}$ its corresponding invertible sheaf. The relative invariants of f are defined as follows:

$$\begin{aligned}\chi_f &= \deg f_*\omega_{X/C} = \chi(\mathcal{O}_X) - (g-1)(g(C)-1), \\ K_f^2 &= K_{X/C}^2 = K_X^2 - 8(g-1)(g(C)-1), \\ e_f &= \chi_{\text{top}}(X) - 4(g-1)(g(C)-1) = \sum_{i=1}^s (\chi_{\text{top}}(F_i) - (2-2g)).\end{aligned}$$

These invariants are nonnegative, and $K_f^2 = 0$ (equivalently, $\chi_f = 0$) if and only if f is locally trivial. $e_f = 0$ iff f is smooth. Let

$$e_{F_i} = \chi_{\text{top}}(F_i) - (2-2g).$$

If F_i is semistable, then e_{F_i} is equal to the number of nodes of F_i .

Arakelov [1] and Beauville [5] proved that K_f is a nef divisor and the curves E with $EK_f = 0$ are those (-2) -curves in the fibers. The map defined by $|nK_f|$ for large n is the contraction morphism $\sigma : X \rightarrow X^\#$ of the vertical (-2) -curves (cf. [12]), we get the relative canonical model $f^\# : X^\# \rightarrow C$ of f .

In what follows, we assume that $f : X \rightarrow C$ is semistable with s singular fibers. Denote by q a singular point of $X^\#$. Then $(X^\#, q)$ is a rational double point of type A_{μ_q} , here μ_q is the number of (-2) -curves in X over q . We also denote by q the singular point of the fibers on the smooth part of $X^\#$, in this case $\mu_q = 0$. Then

$$e_f = \sum_q (\mu_q + 1)$$

is the number of nodes in the singular fibers. For convenience, we let

$$r_f := \sum_q \frac{1}{\mu_q + 1}.$$

It is obvious that

$$r_f \leq e_f. \tag{1.1}$$

In [11], we proved the strict canonical class inequality when $s \neq 0$:

$$K_f^2 < (2g-2)(2g(C)-2+s).$$

The inequality follows from the following

$$K_f^2 - (2g-2)(2g(C)-2+s) \leq -\frac{(2g-2)s}{e} + \frac{3 \cdot r_f}{e^2}, \tag{1.2}$$

where $e \geq 2$ is any integer (see [11], p.594, (5)). Equivalently,

$$r_f \geq \frac{1}{3}e^2 \left(K_X^2 - (2g-2) \left(6g(C) - 6 + s - \frac{s}{e} \right) \right). \tag{1.3}$$

Lemma 1.1. *Let q_1, \dots, q_r be the points such that $\mu_{q_i} \neq 0$. Let $\ell' = \sum_{\mu_q \neq 0} \mu_q$ be the number of (-2) -curves contained in the fibers of f . Then*

$$r_f \leq e_f - \ell' - \frac{r}{2}. \tag{1.4}$$

Proof. Note that the number of points q such that $\mu_q = 0$ is $e_f - \sum_i (\mu_{q_i} + 1)$. Hence

$$\begin{aligned} \sum_q \frac{1}{\mu_q + 1} &= e_f - \sum_i (\mu_{q_i} + 1) + \sum_i \frac{1}{\mu_{q_i} + 1} \\ &\leq e_f - \ell' - r + \frac{r}{2} \\ &= e_f - \ell' - \frac{r}{2}. \end{aligned}$$

This completes the proof. \square

Lemma 1.2. $e_{F_i} = g - 1 + c_i - g(\tilde{F}_i)$, where c_i is the number of components of F_i and \tilde{F}_i is the normalization of F_i . So

$$e_f \leq s(g - 1) + \sum_{i=1}^s c_i. \quad (1.5)$$

Proof. Let $\sigma : \tilde{F}_i \rightarrow F_i$ be the normalization of F_i , and Δ be the subscheme of the singular locus of F_i . Then we have

$$0 \rightarrow \mathcal{O}_{F_i} \rightarrow \sigma_* \mathcal{O}_{\tilde{F}_i} \rightarrow \mathcal{O}_{\Delta} \rightarrow 0.$$

Since F_i has only nodes as its singular points, we have $e_{F_i} = \deg \Delta = h^0(\mathcal{O}_{\Delta}) = \chi(\mathcal{O}_{\Delta})$. Thus

$$\begin{aligned} e_{F_i} &= \chi(\mathcal{O}_{\tilde{F}_i}) - \chi(\mathcal{O}_{F_i}) \\ &= c_i - g(\tilde{F}_i) - (1 - g) \\ &\leq g - 1 + c_i. \end{aligned}$$

This completes the proof. \square

Corollary 1.3. Denote by $\ell = \sum_i c_i - \ell'$ the number of curves in the singular fibers different from (-2) -curves. Then we have

$$r_f \leq s(g - 1) + \ell - \frac{r}{2}. \quad (1.6)$$

Proof. This follows from (1.4) and (1.5). \square

Lemma 1.4. If $e^2 > \frac{6g+3}{g-1}$, then

$$K_f^2 \leq \frac{2e(g-1)^2((2g(C)-2+s)e-s)}{e^2(g-1)-3(2g+1)}. \quad (1.7)$$

Proof. By Cornalba-Harris-Xiao's inequality [6, 14],

$$K_f^2 \geq \frac{4g-4}{g} \chi_f,$$

we get

$$e_f \leq \frac{2g+1}{g-1} K_f^2.$$

Then from (1.1) and (1.2), we obtain

$$K_f^2 - (2g-2)(2g(C)-2+s) \leq -\frac{(2g-2)s}{e} + \frac{3(2g+1)}{e^2(g-1)} K_f^2,$$

it implies (1.7). \square

2 The proof of Theorem 0.2

In this section, we let $f : X \rightarrow \mathbb{P}^1$ be a relatively minimal fibration of genus $g \geq 2$. If $\kappa(X) \neq -\infty$, we denote by S the unique minimal model of X .

Theorem 2.1. *Assume that f is not locally trivial. Then*

$$K_f^2 \geq \begin{cases} 4g - 4, & \text{if } \kappa(X) = -\infty, \\ 6g - 6, & \text{if } \kappa(X) = 0, \\ 6g - 5, & \text{if } \kappa(X) = 1, \\ 6g - 6 + \frac{1}{2} \left(K_S^2 + \sqrt{K_S^2} \sqrt{K_S^2 + 8g - 8} \right), & \text{if } \kappa(X) = 2. \end{cases}$$

- (1) $K_f^2 = 4g - 4$ if and only if X is the minimal resolution of the singularities of a double covering surface $Z \xrightarrow{\pi} \mathbb{P}^1 \times C$ ramified over a curve of numerical type $2F_1 + (2g + 2 - 4g(C))F_2$, and the fibration f is induced by the first projection $\mathbb{P}^1 \times C \rightarrow \mathbb{P}^1$. Here F_i is a fiber of the i -th projection of $\mathbb{P}^1 \times C$.
- (2) If $\kappa(X) = 0$, then $K_f^2 = 6g - 6$ iff the fibration is induced by a pencil $\Lambda \subset |C|$ on its minimal model S with $C^2 = 2g - 2$ simple base points (including infinitely near base points).
- (3) If $\kappa(X) = 1$, then $K_f^2 = 6g - 5$ iff the fibration is induced by a pencil $\Lambda \subset |C|$ of genus g on its minimal model S with $C^2 = 2g - 3$ simple base points.
- (4) If $\kappa(X) = 2$, then $K_S^2 \geq 1$. $K_f^2 = 6(g - 1) + \frac{1}{2} \left(K_S^2 + \sqrt{K_S^2} \sqrt{K_S^2 + 8g - 8} \right)$ iff the fibration is induced by a pencil $\Lambda \subset |C|$ on S with only simple base points and $C \sim rK_S$. In particular, if $g = 2$, then $K_f^2 \geq 8$, with equality iff $K_S^2 = 1$ and $C \sim K_S$. If $g = 3$, then $K_f^2 \geq 15$.

Proof. (1) Let $A = K_X + F$. We claim that $|2A|$ (resp. $|3A|$) is base point free if $g \geq 3$ (resp. $g = 2$). In particular, A is nef and hence

$$K_f^2 - 4(g - 1) = A^2 \geq 0.$$

Indeed, by Ramanujan's vanishing theorem ([2], p.131), $H^1(-F) = H^0(-F) = 0$. So

$$h^0(A) = h^2(-F) = \chi(-F) = \chi_f > 0.$$

Hence we can assume that A is an effective divisor. Since $AF = 2g - 2 \geq 2$, A admits at least one horizontal component. Hence $AK_f \geq 1$.

Note that $|2A| = |K_X + K_f|$. If $g \geq 3$, $K_f^2 = AK_f + 2(g - 1) \geq 1 + 4 = 5$. Suppose $|2A|$ has a base point p , by Reider's theorem [10], there is a curve E passing through p such that

- (i) $K_f E = 0$ and $E^2 = -1$; or (ii) $K_f E = 1$ and $E^2 = 0$.

On the other hand, $K_f E = K_X E + 2FE \equiv K_X E \equiv E^2 \pmod{2}$. Thus the two cases can not exist. This proves that $|2A|$ has no base point.

Now we consider the case $g = 2$. We first prove that A is nef.

Suppose Γ is an irreducible and reduced curve with $A\Gamma < 0$. It is easy to see that Γ must be a horizontal curve. Hence $(A + F)\Gamma = K_f \Gamma \geq 1$. Then we have $\Gamma F \geq 2$. On the other hand, $|A|$ is non empty, so Γ is the fixed part of $|A|$. Hence $2 = 2g - 2 = AF \geq \Gamma F \geq 2$. We get $\Gamma F = 2$, and $A = \Gamma + E$, where E consists of vertical curves. It implies that $\Gamma^2 = A\Gamma - E\Gamma < 0$. Since $A\Gamma = K_X \Gamma + 2 < 0$, we have $K_X \Gamma \leq -3$. So $p_a(\Gamma) < 0$, impossible. Therefore, A is nef and $A^2 \geq 0$.

In this case, $|3A| = |K_X + 2A + F|$. $L = 2A + F$ is nef and $L^2 = 4A^2 + 4AF \geq 8(g-1) > 4$. If $|3A|$ has a base point p , then there is a curve E passing through p such that

(i) $LE = 0$, $E^2 = -1$; or (ii) $LE = 1$, $E^2 = 0$.

Since A and F are nef, (i) implies $AE = FE = 0$. Note that $A = K_X + F$, we have $K_X E = 0$ and $K_X E \neq E^2$, a contradiction. So case (i) is impossible. In case (ii), we have $AE = 0$ and $FE = 1$, so $K_X E = -1$ which is also impossible since $E^2 = 0$. This proves that $|3A|$ is base point free.

Now we consider the case when $K_f^2 = 4g - 4$, i.e., $A^2 = 0$.

In this case, the base point free linear system $|6A|$ is composed with a fibration $\varphi : X \rightarrow C$ over a smooth curve C . Denote by F' a generic fiber of φ . Then $AF' = 0$. Since $AF = 2g - 2 \geq 2$, F' can not be the fiber of f . So $FF' \geq 1$. From $AF' = 0$, we get $K_X F' = -FF' \leq -1$. Since $F'^2 = 0$, we have $K_X F' = -2$ and $FF' = 2$. So $F' \cong \mathbb{P}^1$. Now we get a generally double cover $\pi : X \rightarrow \mathbb{P}^1 \times C$ defined by $\pi(x) = (f(x), \varphi(x))$. Since the pullback $\pi^* F_2$ of a generic fiber F_2 is still isomorphic to \mathbb{P}^1 , and a generic $\pi^* F_1$ has genus g . By Hurwitz formula, we see that the branch curve B_π of π has numerical type $2F_1 + (2g + 2 - 4g(C))F_2$.

Conversely, if the branch curve B_π has numerical type $2F_1 + (2g + 2 - 4g(C))F_2$, then B_π is nonsingular or admits at most ADE singularities. Thus the canonical resolution is the minimal one, and the fibration induced is relatively minimal. By easy computations ([2], p.183, or [8]), we have $K_X^2 = -4(g-1)$ and hence $K_f^2 = 4g - 4$. This completes the proof of (1).

(2) Suppose that the Kodaira dimension of X is non negative, and let S be the unique minimal model of X obtained by contracting (-1) -curves. So the fibers F of f are contracted to a pencil Λ in $|C|$ on S with base points p_1, \dots, p_m (including infinitely near base points). We consider a generic curve C in Λ , and let n_i be the multiplicity of C at p_i . Then $n_i \geq 1$. Note that $F^2 = 0$, $K_X F = 2g - 2$. One can prove easily that

$$C^2 = F^2 + \sum_{i=1}^m n_i^2 = \sum_{i=1}^m n_i^2. \quad (2.1)$$

$$K_S C = K_X F - \sum_{i=1}^m n_i = 2g - 2 - \sum_{i=1}^m n_i. \quad (2.2)$$

Since S is neither ruled nor rational, K_S is nef and $K_S^2 \geq 0$. Hence $K_S C \geq 0$. Thus $m \leq \sum_i n_i \leq 2g - 2 - K_S C$.

$$K_f^2 = K_X^2 + 8(g-1) = K_S^2 - m + 8(g-1) \geq K_S^2 + K_S C + 6(g-1). \quad (2.3)$$

Hence $K_f^2 \geq 6g - 6$. If $K_f^2 = 6g - 6$, i.e., $K_X^2 = -(2g-2) < 0$, then $K_S C = 0$, $K_S^2 = 0$ and $n_i = 1$ for any i . In particular, $m = K_S^2 - K_X^2 = 2g - 2 > 0$ and $C^2 > 0$. By Hodge index theorem, K_S is numerically trivial, thus $\kappa(X) = 0$. By (2.1), $C^2 = m = 2g - 2$. This completes the proof of (2).

(3) We have proved that if $\kappa(X) \geq 0$ and $K_f^2 = 6g - 6$, then $\kappa(X) = 0$. So if $\kappa(X) = 1$, we obtain $K_f^2 \geq 6g - 5$ and $K_S^2 = 0$. Suppose $K_f^2 = 6g - 5$, i.e., $K_X^2 = -(2g-3) < 0$, then $m = 2g - 3 > 0$ and $C^2 > 0$. As in the proof of (2), $K_S C \neq 0$ (otherwise, by Hodge index theorem, $K_S \sim 0$ and $\kappa(X) = 0$). So $K_S C \geq 1$. From our assumption $K_f^2 = 6g - 5$ and (2.3), we get $K_S C = 1$. Now (2.2) implies that $n_i = 1$ for any i , hence $C^2 = 2g - 3$ and C is smooth. The fibration is induced by a pencil in $|C|$ with $2g - 3$ simple base points.

(4) In the case when $\kappa(X) = 2$, we let $x = CK_S$, $y = C^2$. Then

$$x + y \geq 2g - 2, \quad K_S^2 \cdot y \leq x^2. \quad (2.4)$$

It is easy to prove that

$$x \geq \frac{1}{2} \left(-K_S^2 + \sqrt{K_S^2} \sqrt{K_S^2 + 8g - 8} \right).$$

Now from (2.3), we have

$$K_f^2 \geq 6g - 6 + \frac{1}{2} \left(K_S^2 + \sqrt{K_S^2} \sqrt{K_S^2 + 8g - 8} \right).$$

If the equality holds, then the equalities in (2.4) hold. We obtain easily the desired characterization. \square

Corollary 2.2. *Let $f : X \rightarrow \mathbb{P}^1$ be a relatively minimal semistable fibration of genus $g \geq 2$ with s singular fibers. If f is non trivial and $s = 5$, then X is birationally ruled or rational.*

Proof. By the strict canonical class inequality, we have $K_f^2 < 6g - 6$. The corollary follows from Theorem 2.1. \square

Note that if $s = 5$, then S is either \mathbb{P}^2 or a geometrically ruled surface. So

$$m = K_S^2 - K_X^2 > K_S^2 + 2g - 2 = \begin{cases} 2g + 7, & \text{if } S \cong \mathbb{P}^2, \\ 2g + 6 - 8q(X), & \text{otherwise.} \end{cases}$$

3 The proof of Theorem 0.1 for $s = 6$

In this section, we assume that $f : X \rightarrow \mathbb{P}^1$ is semistable and $s = 6$. We use freely the notations in the previous sections, including those in the proof. Denote by F_1, \dots, F_6 the 6 singular fibers.

We recall the inequalities in Sect. 1 for our f :

$$(A) \quad K_X^2 < 0, \quad \text{by (0.1)}$$

$$(B) \quad r_f \geq \frac{1}{3}e(K_X^2 e + 12g - 12), \quad \text{by (1.3)}$$

$$(C) \quad r_f \leq 6g - 6 + \ell - \frac{r}{2}, \quad \text{by (1.6)}$$

$$(D) \quad K_X^2 \leq 4 \frac{(g-1)^2(2e-3)e}{e^2(g-1) - 6g - 3} - 8(g-1), \quad \text{if } e^2 > \frac{6g+3}{g-1}, \quad \text{by (1.7)}$$

where $r_f = \sum_q \frac{1}{\mu_q + 1}$, and ℓ is the number of non (-2) -curves in the 6 singular fibers.

In what follows, we assume that X is of general type, thus $K_S^2 \geq 1$. Denote respectively by $C_1, \dots, C_6 \subset S$ the image curves of F_1, \dots, F_6 . By (A),

$$m = K_S^2 - K_X^2 \geq 2. \quad (3.1)$$

We see that $K_S C \neq 1$. Otherwise, by Hodge index theorem, $K_S^2 C^2 \leq (K_S C)^2 = 1$, so $C^2 \leq 1$, which implies $m \leq 1$. Hence

$$2 \leq K_S C \leq 2g - 2 - m. \quad (3.2)$$

Theorem 3.1. *If $g \leq 4$, then X is not of general type.*

Proof. We shall rule out the cases $g = 2, 3$ and 4 under the assumption $\kappa(X) = 2$.

Case $g = 2$: (3.2) implies $m = 0$, which contradicts (3.1).

Case $g = 3$: By (2.2), (3.1) and (3.2), $m = K_S C = 2$, $K_S^2 = 1$, $K_X^2 = -1$, $n_1 = n_2 = 1$, hence $C^2 = 2$. By Noether inequality $K_S^2 \geq 2p_g(S) - 4$ and Debarre's inequality [7] $K_S^2 \geq 2p_g(S) \geq 2q(S)$ if $q(S) \geq 1$, we obtain

$$p_g(S) \leq 2, \quad q(S) = 0. \quad (3.3)$$

Note that

$$e_f = 12\chi_f - K_f^2 = 12(p_g(S) + 1) - K_X^2 + 4(g - 1) = 12p_g(S) + 21 \leq 45. \quad (3.4)$$

On the other hand, from (B) for $e = 12$, we get $r_f \geq 48$. This contradicts (1.1): $r_f \leq e_f$.

Case $g = 4$: (D) for $e = 12$ implies that $K_X^2 \leq -8/5$. Thus $K_X^2 \leq -2$ and $m \geq 3$. By (3.2), we see that $m \leq 4$.

If $m = 3$, then $K_S^2 = 1$ and $K_X^2 = -2$, we see that (3.3) holds. As in (3.4), we obtain $e_f \leq 50$. (1.3) for $e = 9$ implies that $r_f \geq 54$, which contradicts $r_f \leq e_f$.

If $m = 4$, then (2.2) and (3.2) imply that $K_S C = 2$, $n_1 = \dots = n_4 = 1$, and $C^2 = 4$. By Hodge index theorem, we can see that $K_S^2 = 1$ and $K_S^2 C^2 = (K_S C)^2$. Hence $C \sim 2K_S$. This means that any (-2) -curve E on S does not pass through any one of the 4 base points p_1, \dots, p_4 because $CE = 2K_S E = 0$. Hence (-2) -curves in C_i must be (-2) -curves in F_i .

Note that $K_S C_i = 2$, so C_i has at most two components different from (-2) -curves. This implies that F_i has at most two components which are neither (-2) -curves nor the exceptional curves of the 4 base points.

On the other hand, among the 4 exceptional curves on X , at least one of them is a horizontal (-1) -curve. If the remaining three exceptional curves are vertical, then at least one of them is a (-2) -curve. Hence the total number ℓ of non (-2) -curves in the 6 singular fibers $\leq 6 \times 2 + 2 = 14$.

Now by (C), $r_f \leq 32$. On the other hand, (B) for $e = 6$ implies that $r_f \geq 36$, a contradiction. \square

Theorem 3.2. *If $s = 6$, $g = 5$ and X is of general type, then the minimal model S of X satisfies*

$$K_S^2 = 1, \quad p_g(S) = 2, \quad q(S) = 0.$$

The fibration comes from a pencil on S with 5 simple base points.

Proof. In this case, (D) for $e = 10$ implies that $K_X^2 \leq -\frac{864}{367} < -2$, thus $K_X^2 \leq -3$ and $m \geq 4$. On the other hand, $m \leq 2g - 2 - 2 = 6$. If $m = 6$, then $C^2 \geq 6$ and $K_S C \leq 2$, by Hodge index theorem, $K_S^2 C^2 \leq (K_S C)^2 \leq 4$, and hence $K_S^2 \leq 0$, a contradiction. Thus $m = 4$ or 5 .

If $m = 4$, then $K_S^2 = 1$, $K_X^2 = -3$, $p_g(S) \leq 2$, $q(S) = 0$, $C^2 \geq 4$, and $K_S C \leq 4$. Thus

$$e_f = 12(p_g + 1) - K_X^2 + 4(g - 1) \leq 55.$$

From (1.3) for $e = 8$, we have

$$r_f \geq \frac{1}{3}e(-3e + 48) = e(16 - e) = 64 > e_f.$$

It contradicts (1.1).

So $m = 5$. By (2.1) and (2.2), we have $C^2 \geq 5$, $K_S C \leq 3$. From (3.2), we get $K_S C \geq 2$. Since $K_S^2 \geq 1$, by Hodge index theorem, $K_S^2 C^2 \leq (K_S C)^2$, we have $K_S C = 3$ and $K_S^2 = 1$. So $K_X^2 = -4$, $p_g \leq 2$ and $q = 0$. By (2.1) and (2.2), we obtain $n_1 = \dots = n_5 = 1$ and $C^2 = 5$. Generic C in the pencil is smooth. Let $e = 6$. Then (B) gives us that $r_f \geq \frac{1}{3}e(-4e + 48) = 48$.

Now we prove that $p_g = 2$. Otherwise $p_g \leq 1$, by (3.4), $e_f \leq 12(1 + 1) - (-4) + 16 = 44 < r_f$, a contradiction. Then we have

$$K_S^2 = 1, \quad p_g = 2, \quad q = 0, \quad m = 5.$$

So $e_f = \sum_q (\mu_q + 1) = 56$, $r_f = \sum_q \frac{1}{\mu_q + 1} \geq 48$.

Let $m_i = \#\{q \mid \mu_q = i\}$. Then

$$\begin{aligned} e_f &= m_0 + 2m_1 + 3m_2 + \cdots = 56, \\ r_f &= m_0 + \frac{1}{2}m_1 + \frac{1}{3}m_2 + \cdots \geq 48. \end{aligned}$$

We see that $e_f - m_0 \geq 4(r_f - m_0)$, hence $m_0 \geq 46$. If $m_0 = 46$, one can prove that $m_1 = 5$ and $m_i = 0$ for $i > 1$. So there are at least 46 points q such that $\mu_q = 0$, and there are at most $(56 - 46)/2 = 5$ points q with $\mu_q \neq 0$. By (1.4) we get

$$\ell' + \frac{r}{2} \leq 8.$$

It implies that $r \leq 5$ and $\ell' \leq 7$. □

4 Two examples

Note that for any two points p_1, p_2 on $\mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$, there exists a cyclic cover $\varphi_{p_1, p_2} : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ ramified exactly over p_1 and p_2 . For example, $\tau_n : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ defined by $x \mapsto x^n$ is totally ramified over 0 and ∞ . Now we are going to construct some covers of \mathbb{P}^1 .

Let $\pi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be the double cover ramified over 1 and ∞ , and let $\psi_e : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be the cyclic cover of degree e totally ramified over the two points $\pi^{-1}(0)$. Then the composition

$$\varphi_{2e} : \mathbb{P}^1 \xrightarrow{\psi_e} \mathbb{P}^1 \xrightarrow{\pi} \mathbb{P}^1$$

is a covering of degree $2e$ ramified uniformly over 0, 1, ∞ . The map φ_{2e} is the quotient map of \mathbb{P}^1 for the standard action of the dihedral group of order $2e$ on \mathbb{P}^1 . The cover has two ramification points over 0 with ramification index e , and e ramification points over 1 (resp. ∞) with index 2. Now we consider the fiber product of $\tau_n : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ and $\varphi_{2e} : \mathbb{P}^1 \rightarrow \mathbb{P}^1$, we get a curve

$$\Gamma = \{(x, y) \in \mathbb{P}^1 \times \mathbb{P}^1 \mid \tau_n(x) = \varphi_{2e}(y)\}.$$

This curve is of type $(2e, n)$. The first projection $p_1 : \Gamma \rightarrow \mathbb{P}^1$ can be viewed as the pullback of φ_{2e} under the base change τ_n . So the set Σ of critical points of p_1 is $\tau_n^{-1}\{0, 1, \infty\}$. Σ contains $n + 2$ points. In fact, Γ is locally defined by $\varphi_{2e}(y) = \tau_n(x)$. Thus we can see that Γ admits two singular points defined by $y^e = x^n$ over $\tau_n^{-1}(0)$ and e singular points of type $y^2 = x^n$ over $\tau_n^{-1}(\infty)$. p_1 has e simple ramification points over each point of $\tau_n^{-1}(1)$.

$$\begin{array}{ccc} \Gamma & \subset & \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{\text{pr}_2} \mathbb{P}^1 \\ p_1 \searrow & \text{pr}_1 \downarrow & 2e:1 \downarrow \varphi_{2e} \\ & \mathbb{P}^1 & \xrightarrow[\tau_n]{n:1} \mathbb{P}^1 \end{array}$$

Let $e = n = 4$. Then Γ is of type $(8, 4)$. Let X be the minimal resolution of the double cover $\Sigma \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ ramified over Γ . X can be obtained by the canonical resolution $\sigma : Y \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$, where σ is the blowing-up of $\mathbb{P}^1 \times \mathbb{P}^1$ at the two singular points of Γ over $\tau_n^{-1}(0)$. Denote by E_1 and E_2 the two exceptional curves, and by $\bar{\Gamma}$ the strict transform of Γ on Y . Then $\bar{\Gamma} \equiv \sigma((8, 4)) - 4E_1 - 4E_2 = 2\bar{\delta}$, where $\bar{\delta} = \sigma^*((4, 2)) - 2E_1 - 2E_2$. Let $\bar{\pi} : \bar{X} \rightarrow Y$ be the double cover ramified over $\bar{\Gamma}$. Note that $\bar{\Gamma}$ admits only ADE singularities, so \bar{X} admits at most rational double points, and X is the minimal resolution of \bar{X} . Now by the formulas ([2], p.183), we have

$$K_{\bar{X}} = \bar{\pi}^*(K_Y + \bar{\delta}) = \bar{\pi}^*(\sigma^*(2, 0) - E_1 - E_2).$$

Denote by C_1 and C_2 the two horizontal sections of $\text{pr}_1 : \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ passing through the two points blown up by σ , and by \bar{C}_i their strict transforms on Y . Since \bar{C}_1 and \bar{C}_2 are disjoint with $\bar{\Gamma}$, it is easy to see that their pullback on \bar{X} are 4 (-1) -curves. Note that

$$\bar{C}_1 + \bar{C}_2 = \sigma^*(2, 0) - E_1 - E_2,$$

so $K_{\bar{X}} = \bar{\pi}^*(\bar{C}_1 + \bar{C}_2)$ is the sum of the 4 (-1) -curves. Let S be the surfaces obtained by contracting the 4 horizontal (-1) -curves. Then we see that $K_S \equiv 0$. By easy computation, we have

$$\begin{aligned} \chi(\mathcal{O}_S) &= \chi(\mathcal{O}_{\bar{X}}) \\ &= 2\chi(\mathcal{O}_Y) + \frac{1}{2}(\bar{\delta}^2 + K_Y \bar{\delta}) \\ &= 2 + 0 = 2. \end{aligned}$$

So S is a $K3$ surface. The first projection of $\mathbb{P}^1 \times \mathbb{P}^1$ induces a semistable fibration $f : X \rightarrow \mathbb{P}^1$ of genus 3 with $n+2 = 6$ singular fibers. Because $K_X^2 = -4$, $K_f^2 = 12 = 6g - 6$.

Similarly, let $e = n = 5$, and let F_∞ be the vertical fiber of $\mathbb{P}^1 \times \mathbb{P}^1$ passing through the two singular points of type $x^5 = y^5$. Then consider the double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ ramified over $F_\infty + \Gamma$. The canonical resolution is similar to the above case. We have $\bar{\delta} \equiv \sigma^*((5, 3)) - 3E_1 - 3E_2$ and

$$\begin{aligned} K_Y + \bar{\delta} &= \sigma^*((3, 1)) - 2E_1 - 2E_2 \\ &= \sigma^*((1, 1)) - E_1 - E_1 + \bar{C}_1 + \bar{C}_2 \\ &= \sigma^*((1, 0)) + \bar{F}_\infty + \bar{C}_1 + \bar{C}_2, \end{aligned}$$

where $\bar{F}_\infty \equiv \sigma^*((0, 1)) - E_1 - E_2$ is the strict transform of F_∞ on Y . The pullback of $\bar{F}_\infty + \bar{C}_1 + \bar{C}_2$ on \bar{X} consists 5 (-1) -curves, the one from \bar{F}_∞ is a vertical (-1) -curve. After blowing-down this vertical (-1) -curve, we get a relatively minimal semistable fibration $f : X \rightarrow \mathbb{P}^1$ of genus $g = 4$ with $n+2 = 7$ singular fibers. We see easily that $|K_S|$ is a genus 2 pencil with one simple base point. The pencil is in fact induced by pr_2 . So X is of general type. By easy computation, we have $\chi(\mathcal{O}_S) = 3$, hence

$$K_S^2 = 1, \quad p_g(S) = 2, \quad q(S) = 0.$$

In this case, $K_X^2 = -3$ and $K_f^2 = 6g - 3$.

Acknowledgment. The first author would like to thank Prof. H. Esnault and Prof. E. Viehweg for their hospitality and useful discussions with Viehweg while he is visiting Universität Essen. The third author would like to thank Prof. A. Beauville for useful comments. We all thank the referee for pointing out a mistake and providing us many valuable suggestions in rewriting the manuscript which make the paper readable.

References

- [1] S. Ju. Arakelov: *Families of algebraic curves with fixed degeneracy*, Math. USSR Izvestija **5** (1971), 1277–1302
- [2] W. Barth, C. Peters, A. Van de Ven: *Compact Complex Surfaces*, Springer Verlag 1984
- [3] A. Beauville: *Le nombre minimum de fibres singulières d’une courbe stable sur \mathbb{P}^1* , in: Séminaire sur les pinceaux de courbes de genre au moins deux (L. Szpiro, ed.), Astérisque **86** (1981), 97–108

- [4] A. Beauville: *Les familles stables de courbes elliptiques sur \mathbb{P}^1 admettant quatre fibres singulières*, C.R. Acad. Sci. Paris **294** (1982), 657–660
- [5] A. Beauville: *L'inégalité $p_g \geq 2q - 4$ pour les surfaces de type général*, Appendice à O. Debarre: *Inégalités numériques pour les surfaces de type général*, Bull. Soc. Math. France **110** (1982), no. 3, 319–346
- [6] M. Cornalba, J. Harris: *Divisor classes associated to families of stable varieties, with applications to the moduli space of curves*, Ann. Sci. École Norm. Sup. (4) **21** (1988) no. 3, 455–475
- [7] O. Debarre: *Inégalités numériques pour les surfaces de type général*, Bull. Soc. Math. France **110** (1982), no. 3, 319–346
- [8] E. Horikawa: *Algebraic surfaces of general type with small c_1^2 . V* J. Fac. Sci. Univ. Tokyo Sect. IA Math. **28** (1981), no. 3, 745–755
- [9] K. Liu: *Geometric height inequalities*, Math. Res. Lett. **3** (1996) no. 5, 693–702
- [10] I. Reider: *Vector bundles of rank 2 and linear systems on algebraic surfaces* Ann. of Math. **127** (1988), 309–316
- [11] S.-L. Tan: *The minimal number of singular fibers of a semistable curve over \mathbb{P}^1* , J. Algebraic Geometry **4** (1995), 591–596
- [12] S.-L. Tan: *Effective behavior of multiple linear systems*, Asian J. of Math. **7** (2003) no. 4
- [13] E. Viehweg: *Positivity of direct image sheaves and applications to families of higher dimensional manifolds*, in: School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000) 249–284 ICTP Lect. Notes, 6, Int. Cent. Theoret. Phys., Trieste, 2001
- [14] G. Xiao: *Fibered algebraic surfaces with low slope*, Math. Ann. **276** (1987) no. 3, 449–466